## Stephen Kelso

## dental implants | oral surgery

Extraction of teeth is an irreversible process and whether routine or difficult is a surgical procedure. As in any surgery, there are some risks. They include, but are not limited to

- 1. Swelling and or bruising and discomfort in the surgery area.
- 2. Stretching of the corners of the mouth resulting in cracking and bruising.
- 3. Possible infection requiring further treatment.
- 4. Dry socket jaw pain beginning a few days after surgery, usually requiring additional care, it is more common in lower extractions, especially wisdom teeth.
- 5. Possible damage to adjacent teeth, especially those with large fillings or caps.
- 6. Numbness or altered sensation in the teeth, lip, tongue and chin, due to the closeness of tooth roots (especially wisdom teeth) to the nerves which can be bruised or injured. Sensation most often returns to normal, but in rare cases, the loss may be permanent.
- 7. Trismus limited jaw opening due to inflammation or swelling, most common after wisdom tooth removal. Sometimes it is the result of jaw joint discomfort (TMJ), especially when TMJ disease and symptoms already exist.
- 8. Bleeding significant bleeding is not common, but persistent oozing can be expected for several hours.
- 9. Sharp ridges or bone splinters may form later at the edge of the socket. These may require another surgery to smooth or remove them.
- 10. Incomplete removal of tooth fragments to avoid injury to vital structures such as nerves or sinuses, sometimes small root tips may be left in place.
- 11. Sinus involvement: The roots of upper back teeth are often close to the sinus and sometimes a piece of root can be displaced into the sinus, or an opening may occur into the mouth which may require additional care.

| (dissolve) without intervention.   |
|--|
| Most procedures are routine and serious complications are not expected. Those, which do occur, are most often minor and can be treated.                                      |
| Teeth to be removed:   |
|  |
|  |
| I have read and understand the above, and had my questions answered. I recognise there can be no guarantee as to the outcome of treatment, and I give my consent to surgery. |
| Patient or legal guardian  |
| Date   |
|  |
| Stephen Kelso BDS MFDS RCPS  |
| Date   |
|  |
|  |
|  |
|  |
|  |

12. Following a surgical procedure usually sutures will be placed. These will resorb